



The emerging role of trogocytosis in the evasion of cancers and parasitic protists from immune cells

Kumiko Nakada-Tsukui¹, Tomoyoshi Nozaki²

¹Department of Parasitology, National Institute of Infectious Diseases, Tokyo, Japan; ²Department of Biomedical Chemistry, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan

Correspondence to: Kumiko Nakada-Tsukui. Department of Parasitology, National Institute of Infectious Diseases, Toyama 1-23-1, Shinjuku-ku, Tokyo 162-8640, Japan. Email: kumiko@nih.go.jp; Tomoyoshi Nozaki. Department of Biomedical Chemistry, Graduate School of Medicine, The University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-8654, Japan. Email: nozaki@m.u-tokyo.ac.jp.

Provenance: This is an invited article commissioned by the Executive Editor-in-Chief Dr. Hualin Sun (Jiangsu Key Laboratory of Neuroregeneration, Nantong University, Nantong, China).

Comment on: Hamieh M, Dobrin A, Cabriolu A, *et al.* CAR T cell trogocytosis and cooperative killing regulate tumour antigen escape. *Nature* 2019;568:112-6.

Received: 25 January 2020; Accepted: 20 February 2020; Published: 31 March 2020.

doi: 10.21037/biotarget.2020.03.02

View this article at: <http://dx.doi.org/10.21037/biotarget.2020.03.02>

The word of “trogocytosis” started to be used in immunology since 2003. The word “trog” means “nibble” or “chew” in Greek, and trogocytosis represents a mode of internalization of a part of a neighboring cell, which is a distinct manner from conventional “phago”cytosis, meaning, as we understood, “eat” with one bite. After the report in 1972 (1), in which transfer of major histocompatibility complex II (MHC class II) from B to T cells was observed, a similar mode of transfer of surface molecules has been reported for a variety of immune cells such as T, B, natural killer (NK), and dendritic cells (2-5). However, it remained elusive whether or not and, if so, how surface molecules such as transmembrane receptors were transferred to other cells. The hypothetical idea of the nibbling of surface membranes was proposed by Hudrisier and colleagues in 2001 (6). In this study, it was shown that surface lipids and proteins were acquired with peptide-loaded MHC and T cell receptor and the phenomenon was named “trogocytosis” (3,6). After their findings, the concept has been generally accepted that surface receptors are transferred between immune cells via trogocytosis. In the recent publication by Hamieh and colleagues, a new role of trogocytosis in immune evasion has been demonstrated, in which tumor cells escape from killing during immune therapy by chimeric antigen receptors (CARs) (7).

CARs are artificially designed receptors that harbor

one extracellular single chain variable fragment (scFv) domain that recognizes target cells and one to three cytosolic intercellular signaling domains derived from CD3 ζ , costimulatory proteins such as CD28, and 4-1BB. Recognition of the target tumor cells by CAR-expressing T cells via the scFv domain causes activation of the signaling domains, which in turn elicit downstream effector activity, such as secretion of perforin, granzyme, and stimulation of Fas, to eliminate tumor cells. Adoptive transfer of CAR-expressing T cells showed impressive efficacy to B cell malignancies and was considered as a promising therapeutic measure (8). However, even though complete remission of tumor cells was observed, relapse was reported to occur because of loss or reduction of the tumor antigen on the leukemia cells and exhaustion and elimination of CAR T cells (9,10). Hamieh and colleagues showed that CD19, a B cell antigen, on the acute lymphoblastic leukaemia (ALL) cells (NALM6) was taken up by CAR T cells via trogocytosis and it is caused reduction of the tumor antigen (7). Subsequently, trogocytosed CD19 in the CAR T cell was transferred onto the CAR T cell surface. This B cell antigen expressing CAR T cell was recognized as malignant B cells and killed by other CAR T cells and it is called T cell fratricide (meaning “murdering one’s brother”) and eventually exhaustion of the CAR T cells (7). Although other mechanisms were reported for immune evasion of

tumors from CAR T cells, such as antigen loss on the tumor cells caused by genetic mutation, alternative splicing, epitope shedding, or lineage switching (11), regulation of trogocytosis could be one of the potential therapeutic strategies.

Trogocytosis is not restricted to mammals. There were a number of reports that described trogocytosis in a wide range of eukaryotes from protists (synonymous to protozoa) to mammals (12). As early as in 1979, Brown and colleagues demonstrated by electron microscopy that the cytopathic mechanism, which was also referred as “trogocytosis”, of the free-living amoeba, *Naegleria fowleri*, by which the amoeba nibbled mouse embryonic cells (13). Similar trogocytosis-like ingestion of the target cells and trogocytosis-mediated cytopathic mechanisms have been reported from other free living and parasitic amoebas, such as *Acanthamoeba*, *Dictyostelium*, and *Entamoeba* (12,14). In addition, in the multicellular organisms, trogocytosis has also been reported to be involved in retinal development in *Drosophila* (15), primordial germ cell development in *Caenorhabditis elegans* (16), remodeling of synapses in *Mus musculus* (17), communication between mesenchymal stroma/stem-like cells and cancer cells in *Homo sapiens* (18), and spread of an intercellular pathogen *Francisella tularensis* between macrophages (19). In contrast to the cytopathic consequence of trogocytosis by protists, trogocytosis by multicellular organisms usually does not lead to killing of target cells. The concept is now well accepted that trogocytosis in multicellular organisms in general serves for intercellular communications. However, there are several notable exceptions. Trogocytosis was implicated in cytopathy by neutrophil and macrophages against cancer cells and the human genital parasitic protist, *Trichomonas vaginalis* (20-23).

A series of the reports on trogocytosis in various organisms suggest the sporadic existence of trogocytosis at least among the supergroup Unikonta (containing Opisthokonta and Amoebozoa) of eukaryotes. Does trogocytosis in protists and multicellular organisms differ in the context of mechanisms and functions? A recent study on trogocytosis in the human intestinal protist, *E. histolytica*, the causative agent of amebiasis, demonstrated that amoebic trogocytosis may also serve for immune evasion (24). It was shown that an *E. histolytica* trophozoite which had trogocytosed Jurkat T cell was decorated with Jurkat T cell-derived membrane proteins such as class I MHC (24). *E. histolytica* trophozoites are exposed to complement during invasion and thus, protection from

complement is indispensable for survival. This finding suggests trogocytosis in protists also serves as intercellular communication mechanism and way of decorating their surface with molecules stole from their neighbors.

To improve CAR T cell therapy, one potential approach will be regulating trogocytosis of CAR T cells. To this end, understand the molecular mechanisms is indispensable. Unfortunately, molecular basis of trogocytosis is not well clarified except for trogocytosis specific AGC family kinase in *E. histolytica* (25), most of the report showing molecules shared with phagocytosis (26). Studies of molecular basis of trogocytosis in model organisms and protists will provide novel drug target.

Acknowledgments

Funding: This work was supported in part by Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan to T Nozaki (JP17K19416, and JP18H02650) and K Nakada-Tsukui (JP19H03463), Research Program on Emerging and Re-emerging Infectious Diseases from Japan Agency for Medical Research and Development (AMED) to T Nozaki (JP18fk0108046) and K Nakada-Tsukui (JP19fk0108049j1103).

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

References

1. Cone RE, Sprent J, Marchalonis JJ. Antigen-binding specificity of isolated cell-surface immunoglobulin from thymus cells activated to histocompatibility antigens. Proc Natl Acad Sci U S A 1972;69:2556-60.
2. Huang JF, Yang Y, Sepulveda H, et al. TCR-Mediated internalization of peptide-MHC complexes acquired by T cells. Science 1999;286:952-4.
3. Joly E, Hudrisier D. What is trogocytosis and what is its purpose? Nat Immunol 2003;4:815.
4. Wakim LM, Bevan MJ. Cross-dressed dendritic cells

- drive memory CD8+ T-cell activation after viral infection. *Nature* 2011;471:629-32.
5. Aucher A, Magdeleine E, Joly E, et al. Capture of plasma membrane fragments from target cells by trogocytosis requires signaling in T cells but not B cells. *Blood* 2008;111:5621-8.
 6. Hudrisier D, Riond J, Mazarguil H, et al. Cutting edge: CTLs rapidly capture membrane fragments from target cells in a TCR signaling-dependent manner. *J Immunol* 2001;166:3645-9.
 7. Hamieh M, Dobrin A, Cabriolu A, et al. CAR T cell trogocytosis and cooperative killing regulate tumour antigen escape. *Nature* 2019;568:112-6.
 8. Brudno JN, Kochenderfer JN. Chimeric antigen receptor T-cell therapies for lymphoma. *Nat Rev Clin Oncol* 2018;15:31-46.
 9. Fry TJ, Shah NN, Orentas RJ, et al. CD22-targeted CAR T cells induce remission in B-ALL that is naive or resistant to CD19-targeted CAR immunotherapy. *Nat Med* 2018;24:20-8.
 10. Caumartin J, Favier B, Daouya M, et al. Trogocytosis-based generation of suppressive NK cells. *EMBO J* 2007;26:1423-33.
 11. Zheng S, Asnani M, Thomas-Tikhonenko A. Escape From ALL-CARTaz: Leukemia Immunoediting in the Age of Chimeric Antigen Receptors. *Cancer J* 2019;25:217-22.
 12. Ralston KS. Taking a bite: Amoebic trogocytosis in *Entamoeba histolytica* and beyond. *Curr Opin Microbiol* 2015;28:26-35.
 13. Brown T. Observations by immunofluorescence microscopy and electron microscopy on the cytopathogenicity of *Naegleria fowleri* in mouse embryo-cell cultures. *J Med Microbiol* 1979;12:363-71.
 14. Ralston KS, Solga MD, Mackey-Lawrence NM, et al. Trogocytosis by *Entamoeba histolytica* contributes to cell killing and tissue invasion. *Nature* 2014;508:526-30.
 15. Cagan RL, Krämer H, Hart AC, et al. The bride of sevenless and sevenless interaction: internalization of a transmembrane ligand. *Cell* 1992;69:393-9.
 16. Abdu Y, Maniscalco C, Heddleston JM, et al. Developmentally programmed germ cell remodelling by endodermal cell cannibalism. *Nat Cell Biol* 2016;18:1302-10.
 17. Weinhard L, di Bartolomei G, Bolasco G, et al. Microglia remodel synapses by presynaptic trogocytosis and spine head filopodia induction. *Nat Commun* 2018;9:1228.
 18. Melzer C, von der Ohe J, Hass R. Concise Review: Crosstalk of Mesenchymal Stroma/Stem-Like Cells with Cancer Cells Provides Therapeutic Potential. *Stem Cells* 2018;36:951-68.
 19. Steele SP, Chamberlain Z, Park J, et al. *Francisella tularensis* enters a double membraned compartment following cell-cell transfer. *Elife* 2019. doi: 10.7554/eLife.45252.
 20. Matlung HL, Babes L, Zhao XW, et al. Neutrophils Kill Antibody-Opsonized Cancer Cells by Trogocytosis. *Cell Rep* 2018;23:3946-59.e6.
 21. Valgardsdottir R, Cattaneo I, Klein C, et al. Human neutrophils mediate trogocytosis rather than phagocytosis of CLL B cells opsonized with anti-CD20 antibodies. *Blood* 2017;129:2636-44.
 22. Mercer F, Ng SH, Brown TM, et al. Neutrophils kill the parasite *Trichomonas vaginalis* using trogocytosis. *PLoS Biol* 2018;16:e2003885.
 23. Velmurugan R, Challa DK, Ram S, et al. Macrophage-Mediated Trogocytosis Leads to Death of Antibody-Opsonized Tumor Cells. *Mol Cancer Ther* 2016;15:1879-89.
 24. Miller HW, Suleiman RL, Ralston KS. Trogocytosis by *Entamoeba histolytica* Mediates Acquisition and Display of Human Cell Membrane Proteins and Evasion of Lysis by Human Serum. *MBio* 2019. doi: 10.1128/mBio.00068-19.
 25. Somlata, Nakada-Tsukui K, Nozaki T. AGC family kinase 1 participates in trogocytosis but not in phagocytosis in *Entamoeba histolytica*. *Nat Commun* 2017;8:101.
 26. Martínez-Martín N, Fernández-Arenas E, Cemerski S, et al. T cell receptor internalization from the immunological synapse is mediated by TC21 and RhoG GTPase-dependent phagocytosis. *Immunity* 2011;35:208-22.

doi: 10.21037/biotarget.2020.03.02

Cite this article as: Nakada-Tsukui K, Nozaki T. The emerging role of trogocytosis in the evasion of cancers and parasitic protists from immune cells. *Biotarget* 2020;4:4.